



PPAR- β Antibody

E11-0556B**Catalog Number:** E11-0556B**Concentration:** 1mg/ml**Swiss-Prot No.:** Q15648

Other Names: ARC205, Activator-recruited cofactor 205 kDa component, DRIP205, PBP, PPAR-binding protein, PPRB, Peroxisome proliferator-activated receptor binding protein, TRIP2, Thyroid hormone receptor associated protein 220, Thyroid hormone receptor-associated protein complex 220 kDa component, Thyroid receptor interacting protein 2, Trap220, Vitamin D receptor-interacting protein complex component DRIP205, p53 regulatory protein RB18A

Storage/Stability: Store at -20°C/1 year

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Immunogen: The antiserum was produced against synthesized peptide derived from Internal of human PPAR- β .

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Specificity: PPAR- β antibody detects endogenous levels of total PPAR- β protein.

Reactivity: Human (Identities = 100%, Positives = 100%);
Mouse (Identities = 92%, Positives = 100%)


Applications:WB: 1:500~1:2000 IF: 1:100~1:500
ELISA: 1:10000

References:

Hiroshi Mano, J. Biol. Chem., Mar 2000; 275: 8126.

Ken-ichiro Otsuyama, Blood (ASH Annual Meeting Abstracts), Nov 2006; 108: 5085.

Liliane Michalik, J. Cell Biol., Aug 2001; 154: 799.



Peptide

-

+

Immunofluorescence analysis of HeLa cells, using PPAR- β antibody.

For Research Use Only