RAC3 (NM_005052) Human Over-expression Lysate

E45H18839-

Catalog Number:E45H18839-2

NCBI Gene ID:5881

Calculated MW:21.4 kDa

Storage: Upon receiving, store the sample at -80°C. After dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2 x SDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -80°C.

Cite ID#:EN5540119

Type: Human Lysates

Research Field: Adherens junction, Axon guidance, B cell receptor signaling pathway, Colorectal cancer, Fc epsilon RI signaling pathway, Focal adhesion, MAPK signaling pathway, Natural killer cell mediated cytotoxicity, Pancreatic cancer, Pathways in cancer, Regulation of actin cytoskeleton, VEGF signaling pathway, Viral myocarditis, Wnt signaling pathway

Description: Transient overexpression lysate of ras-related C3 botulinum toxin substrate 3 (rho family, small GTP binding protein Rac3) (RAC3)

Protein length: GoldORF Clone [GH33278V]

Expression Host:HEK293T

Species: Human

Tag:C-Myc/DDK

Detection principle: Detection Antibody: Clone LBI4C5, Anti-DDK (FLAG) monoclonal antibody ([AMM08808V])

Kit components:1 vial of gene specific transient over-expression cell lysate in RIPA buffer 1 vial of whole HEK293T cell lysate in RIPA buffer 1 vial of 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% glycerol, 0.002% Bromophenol blue, 100mM DTT)

Gene symbol:RAC3

mRNA Refseq:NM_005052

Protein Refseq:NP_005043

Chromosome location:17q25.3

Pathway: Adherens junction, Axon guidance, B cell receptor signaling pathway, Colorectal cancer, Fc epsilon RI signaling

For Research Use Only

pathway, Focal adhesion, MAPK signaling pathway, Natural killer cell mediated cytotoxicity, Pancreatic cancer, Pathways in cancer, Regulation of actin cytoskeleton, VEGF signaling pathway, Viral myocarditis, Wnt signaling pathway

Function: Druggable Genome

Form: Liquid

Preparation:HEK293T cells in 10-cm dishes were transiently transfected with Leading Biology Transfection Reagent (TF002V) and 5ug Leading Biology GoldORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1 x Proteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.

Synonyms:ras-related C3 botulinum toxin substrate 3 (rho family, small GTP binding protein Rac3); rho family, small GTP binding protein Rac3